arXiv:1304.7008v1 [cond-mat.stat-mech] 25 Apr 2013

Memory as order-parameter on a Cayley tree

Javier M. Magán

Auditya Sharma

International Institute of Physics - Federal University of Rio Grande do Norte, Natal, RN, Brazil

Abstract. We provide a concrete and systematic connection between the statistical physics of the Ising ferromagnet on a Cayley tree, which is an old solved problem, and the study of memory in exponentially expanding spaces, which is of current interest. We provide the order parameters which clearly signal the (known) phase transitions. The correlators are found in a new simple way, which may be of interest for other tree like systems, and which allows the computation of higher order expectation values of the order parameter. Numerical Monte Carlo simulations are used to make transparent the existence of the two different transition temperatures.

1. Introduction

The objective of this paper is to make a concrete and transparent connection between two separate fields of study that are interested in the same problem [1, 2, 3, 4, 5, 6] and to bring to the fore old work, which is of very current interest, in modern language from a contemporary perspective.

Our motivation to revisit an old statistical physics problem came from recent works showing the natural appearance of tree like structures in certain gravity phenomena, like bubble nucleation in eternal inflation [7] or in the near horizon region of black holes [8]. The Ising model in the Cayley tree is complicated enough that some interesting questions that arise here can be meaningfully posed, but still simple enough to answer all of them in a clear way.

In particular, in this article we develop the connection between questions related to memory, which have their root in the branching Markov process approach, and which were raised in a different framework by Ref.[9], and the more usual static approach, dealing with partition functions and correlators. Building on old knowledge of this system and with the aid of new techniques to compute certain correlation functions in the Cayley tree, we obtain a clear and simple way the two known critical temperatures, the so called Bethe-Peierels [1] and spin glass [10] critical temperatures. Furthermore, we point out that the expectation value of the product of the spin at the root and the total magnetization at the boundary $\langle\sigma_0 M_n\rangle$ is the appropriate order parameter to signal T_{BP}, and $\frac{\langle\sigma_0 M_n\rangle}{\Omega}$ where $\Omega = \sqrt{\langle(\sigma_0 M_n)^2\rangle - \langle\sigma_0 M_n\rangle^2}$ is the RMS deviation, is the order parameter for T_{SG} - facts that appear to be not widely known. We also provide a rigorous derivation of the Markov transition matrix from the static approach. Also, we point out that this conceptual framework can be easily extended to the spin-glass and Heisenberg-type variants of the Hamiltonian - a discussion of some consequences is included.

Monte Carlo simulations on these type of systems in which the boundary is of the same size as the system itself have always been problematic, due to subtleties with the thermodynamic limit. With the help of this new analytic understanding we are able to perform Monte Carlo simulations, showing clearly the two transition temperatures.

We make clear right at the outset that we consider here the Cayley tree and not the Bethe lattice, the difference between which two has been a source of confusion in the literature - look for example at Ostilli [11].

2. The Static Approach: Partition Function

In 1974, Eggarter [1] provided a beautiful way of computing the partition function of the Ising model on the Cayley tree. The method is based on two important properties. The first one is that, generically, the Ising model Hamiltonian is just a sum of 'bonds', i.e, a sum of products of adjacent spins. The second is that the Cayley tree has no loops and, therefore, the bonds provide an equally good basis in which to define the partition



Figure 1. Figure shows a Cayley tree of depth $n = 3$ and coordination number $z = 3$.

function. The advantage is that the Ising Hamiltonian decouples in these variables. More concretely, we start with the following Hamiltonian:

$$\mathcal{H} = -J \sum_{\langle i,j \rangle} \sigma_i \sigma_j, \tag{1}$$

where the sum involves pairs of spins which are adjacent on the tree. Instead of the spin variables, we choose the spin at the root of the tree σ_0, and the bonds $\theta_{ij} = \sigma_i \sigma_j$ to define the configuration of our system. The θ variables can take values ± 1, which make them effectively spin variables too. The Hamiltonian then takes the following simple form

$$\mathcal{H} = -J \sum_{\langle i,j \rangle} \theta_{ij}. \tag{2}$$

We see that the Ising model on the Cayley tree is essentially a sum of independent spins (the bonds) subjected to a constant magnetic field; in the computation of the partition function, the only difference with this dual system is a factor of two coming from the sum over the possible states of the spin at the root. Therefore we arrive at [1]:

$$Z(J)_{Cayley} = 2(2\cosh(\beta J))^{N_b} = 2(2\cosh(\beta J))^{N-1}, \tag{3}$$

where N_b is the number of bonds and N is the number of spins.

With this formalism, namely the θ variables, the computation of the correlators is straightforward. Firstly, odd correlators vanish due to the spin symmetry of the model. For the even case we begin with the two point function. We would like to write the spin product in terms of θ variables. To achieve this, we just need to find the path between the two spins. This path is unique because there are no closed paths in a Cayley tree. Imagine that the path goes over $\sigma_a, \sigma_b, \cdots, \sigma_c$. We can then write the product as

$$\sigma_i \sigma_j = \sigma_i \sigma_a \sigma_a \sigma_b \sigma_b \cdots \sigma_c \sigma_c \sigma_j = \theta_{ia} \theta_{ab} \cdots \theta_{cj}. \tag{4}$$

The important point now is that, as commented before, the θ variables are not coupled to each other, and are thus independent variables obeying the same Hamiltonian. Therefore,

$$\langle\sigma_i\sigma_j\rangle = \langle\theta_{ia}\theta_{ab}\cdots\theta_{cj}\rangle = \langle\theta_{ia}\rangle\langle\theta_{ab}\rangle\cdots\langle\theta_{cj}\rangle = \langle\theta\rangle^{d_{ij}}, \tag{5}$$

where d_{ij} is the discrete natural distance on the tree, and θ is a spin variable obeying the following simple Hamiltonian

$$\mathcal{H}_\theta = -J\theta \tag{6}$$

with partition function

$$z_\theta = 2\cosh(\beta J), \tag{7}$$

and whose one-point function is given by

$$\langle\theta\rangle = \frac{\sum_{\theta=\pm1}\theta e^{\beta J\theta}}{z_\theta} = \frac{e^{\beta J}-e^{-\beta J}}{e^{\beta J}+e^{-\beta J}} = \tanh(\beta J). \tag{8}$$

Therefore we have, with the defintion $a \equiv \tanh(\beta J)$ which will prove useful later,

$$\langle\sigma_i\sigma_j\rangle = \tanh(\beta J)^{d_{ij}} = a^{d_{ij}}. \tag{9}$$

Going to higher order correlation functions is also possible. The above technique allows us to obtain the general result for a $2n$ point function, namely:

$$\langle\sigma_{i_1}\sigma_{i_2}\cdots\sigma_{i_{2n}}\rangle = (\tanh(\beta J))^{[\sum_{\langle AB\rangle} d_{i_A i_B}]_{min}} \tag{10}$$

where $d_{i_A i_B}$ are the distances between pairs and the minimization is carried out over all possible ways in which n-pairs can be formed from $2n$ points. A simple geometric construction to find this minimum is to select n-pairs at random, join all the paths between pairs, and delink any portions on the tree that are traversed an even number of times. The newly formed pairs will give a minimum for the sum.

3. The Dynamic Approach: Evolution of a Markov Chain

Now we consider the type of approach adopted in Refs. [6, 9]. The problem is the following. Consider an Ising spin at time $t = 0$, which can be up or down with probabilites $P_0 = (p_0^{\uparrow}, p_0^{\downarrow})$. At time t_1 this spin gives rise to z spins, each of which then gives rise to $(z-1)$ spins at the next time instant t_2, each of which in turn gives rise to $(z-1)$ spins, and so on. There is an obvious tree structure appearing here, with the time t playing the role of the lattice depth in the past static scenario. Now the dynamics is determined by a Markov rule: the probability distribution of a spin at level t_{n+1} is completely characterized by the one of its parent spin at level t_n. This process defines a Markov chain, in particular a Branching Markov chain, due to the exponential growth of the number of spins at every time step. A Markov chain is defined by its transition matrix. In our ca se, taking into account that the probability ratios should correspond to the ones coming from the Gibbs distribution we have the following transition matrix:

$$G(\sigma,\sigma') = \frac{e^{\beta J\sigma\sigma'}}{2\cosh(\beta J)} = \frac{e^{\beta J\sigma\sigma'}}{z_\theta} \tag{11}$$

The Markov dynamics is then defined as

$$P_{n+m} = G^m P_n. \tag{12}$$

A rigorous technique, which uses the above θ variables, to obtain the transition matrix, is the following (for clarity we define the transition matrix between the spin at the root and one of its descendants but the computation can be done similarly for any pair of spins) :

The joint probability density of the spin at the root and one of its adjacent spins is given by

$$\begin{aligned} P(\sigma_0,\sigma_1) &= \sum_{\sigma\neq\sigma_0,\sigma_1} \frac{e^{-\beta H}}{Z} = \frac{e^{J\beta\theta_{01}}}{Z}[\mathrm{Tr}_\theta\, e^{-J\theta}]^{N_b-1} \\ &= \frac{e^{J\beta\theta_{01}}}{Z} z_\theta^{N_b-1} = \frac{e^{J\beta\theta_{01}}}{2z_\theta} \end{aligned} \tag{13}$$

and the reduced probability for the spin at the root is just (as expected)

$$P(\sigma_0) = \sum_{\sigma\neq\sigma_0} \frac{e^{-\beta H}}{Z} = \frac{\sum_\theta e^{-\beta H}}{\sum_{\sigma,\theta} e^{-\beta H}} = \frac{z_\theta^{N_b}}{2z_\theta^{N_b}} = \frac{1}{2} \tag{14}$$

The transition matrix is, by definition, the conditional probability $P(\sigma_1|\sigma_0)$, and so

$$G(\sigma_1,\sigma_0) = \frac{P(\sigma_0,\sigma_1)}{P(\sigma_0)} = \frac{e^{J\beta\theta_{01}}}{z_\theta} = \frac{e^{\beta J\sigma\sigma'}}{z_\theta}, \tag{15}$$

which is the transition matrix of Eq.11. Now, in order to compute the correlations, we first diagonalize the matrix G. It is easy to check that, with $a \equiv \tanh(\beta J)$,

$$G = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix} \tag{16}$$

and therefore

$$P_{n+m} = \begin{pmatrix} \frac{1}{2}(1+a^n) & \frac{1}{2}(1-a^n) \\ \frac{1}{2}(1-a^n) & \frac{1}{2}(1+a^n) \end{pmatrix} P_m \tag{17}$$

and as expected, no matter the initial distribution, the probability distribution at late times tends to the stationary distribution $P = (1/2, 1/2)$.

The correlator between any two spins is given by

$$\langle\sigma_i\sigma_j\rangle = \sum_{\sigma_i,\sigma_j=\pm1} \sigma_i\sigma_j P(\sigma_i,\sigma_j) \tag{18}$$

where $P(\sigma_i,\sigma_j)$ is the probability of finding σ_i, σ_j. This probability can be found by going to the first common ancestor of the two spins, σ_r. It is given by

$$P(\sigma_i,\sigma_j) = \sum_{\sigma_r=\pm1} P(\sigma_i|\sigma_r)P(\sigma_j|\sigma_r)P_{\sigma_r} \tag{19}$$



Figure 2. Monte Carlo simulation of Cayley trees with coordination number $z = 3$. Figure (a) obtains the transition temperature popularly called Bethe-Peirels T_{BP}. Figure (b) obtains the so-called spin-glass transition temperature [10], T_{SG} deeper down in the ordered phase. Appendix C has some details of the simulation.Appendix D shows why the second-transition temperature carries the name 'spin-glass': quite remarkably, the same transition temperature appears when one considers the $\pm J$ spin glass on a Cayley tree.

Taking into account that individual spins, and in particular the common ancestor, have stationary distributions in the thermal ensemble, and that the conditional probabilities $P(\sigma_i|\sigma_r)$ are just given by the elements of the matrix $G^{d_{ir}}$ we arrive finally at the desired result

$$\langle\sigma_i\sigma_j\rangle = a^{d_{ij}} \tag{20}$$

So the Markov chain, which is a time process, gives the same value for the correlators between the spins as the common static procedures on a thermal tree. This observation may be interesting for the model studied in Ref.[7], which, in this perspective, may be expected to be equal to some statistical model on a thermal tree. Several observations done in that reference become very natural in this perspective (like the p-adic structure of the boundary correlators). We leave these interesting issues for the near future.

4. The two phase transitions

In the Markov chain approach, a very natural question to ask is how correlated some random variable at the n^{th} time-step is with respect to the root spin; in other words, how much *memory* has that random variable about its earliest ancestor. For example, the correlator $\langle\sigma_0\sigma_j\rangle$ dies for large distances, no matter the value of βJ. So, as commented in Ref.[9], there is no memory for these two random variables. We now show that, in fact, the consideration of the correlation between the *sum* of the spins at the n^{th} time-step with respect to the root-spin, i.e. $\langle\sigma_0 M_n\rangle$, is more useful and gives the order-parameter for the so-called 'Bethe-Peirels' transition. Furthermore, we will show that the consideration of the allied quantity $\frac{\langle\sigma_0 M_n\rangle}{\Omega}$ where $\Omega = \sqrt{\langle(\sigma_0 M_n)^2\rangle - \langle\sigma_0 M_n\rangle^2}$ is the RMS deviation, gives the so-called spin glass transition temperature. Appendix A carries a discussion of the motivation behind studying these two quantities from a generic perspective, and in particular its conceptual relation with the mutual information between the two random variables.

Indeed, that the Ising ferromagnet on a Cayley tree has two distinct phase transitions is long known in the statistical physics community - this information is buried in the papers of Eggarter [1], Matsuda [2], and particularly, the papers of J von Heimburg and Thomas [4], and Morita and Horiguchi [3]; however, we believe that our paper shows the correct order-parameters for these two transitions for the first time.

Computing the correlation between the root spin and the sum of the spins at the n^{th} level we obtain:

$$\langle \sigma_0 M_n \rangle = \sum_{i \in n^{th} layer} \langle \sigma_0 \sigma_i \rangle = N_n \tanh(\beta J)^n, \tag{21}$$

where N_n is equal to the number of spins in the n^{th} layer: $N_n = z(z-1)^{n-1}$, where z is the coordination number for the type of Cayley tree shown in Fig.1. Therefore, defining $\gamma \equiv z - 1$, we have

$$\langle \sigma_0 M_n \rangle = (\gamma \tanh(\beta J))^n (1 + \frac{1}{\gamma}) \tag{22}$$

It is clear that the behaviour of $\langle \sigma_0 M_n \rangle$ for 'long times' or, equivalently, in the 'thermodynamic limit' $n \to \infty$, changes completely depending on whether $\gamma \tanh(\beta J)$ is $>$ or $<$ 1. So, $\gamma \tanh(\beta J) = 1$ signals a transition which is understood to be the so-called Bethe-Peirels critical temperature $T_{BP} = \frac{J}{\tanh^{-1}(\frac{1}{\gamma})}$. Fig.2(a) shows that this phase transition can be obtained from Monte Carlo simulations of finite-systems.

For computing the deviation of this random variable we first need to compute the second order moment. It is given by

$$\begin{aligned} \langle (\sigma_0 M_n)^2 \rangle &= \sum_{i,j \in n^{th} layer} \langle \sigma_i \sigma_j \rangle = \sum_{i,j \in n^{th} layer} a^{d_{ij}} \\ &= (\gamma + 1)\gamma^{n-1} + 2 \sum_{i<j} a^{d_{ij}} \end{aligned} \tag{23}$$

The last sum above was done in Ref.[3]. We compute it in a different way. In particular we can derive a recursion relation for the sum, the details of which are provided in Appendix B. We collect here the result:

$$\sum_{i<j} a^{d_{ij}} = N_n \left(\frac{(\gamma - 1)}{2\gamma} \frac{x^{n+1} - x}{x - 1} + \frac{x^n}{2} \right), \tag{24}$$

where $x \equiv \gamma a^2 \equiv \gamma \tanh^2(\beta J)$, and $N_n = (\gamma + 1)\gamma^{n-1}$ is, once again, the number of spins in the n^{th} layer. With the help of Eq.C.1, and some algebra, we can now write down the mean square deviation:

$$\Omega^2 = \langle (\sigma_0 M_n)^2 \rangle - \langle \sigma_0 M_n \rangle^2 = N_n \left(1 + \frac{\gamma - 1}{\gamma} \frac{x^{n+1} - x}{x - 1} - \frac{1}{\gamma} x^n \right) \tag{25}$$

From this expression one can easily check that the ratio $\langle \sigma_0 M_n \rangle / \Omega$ defines two phases delimited by the second critical temperature, which is given by $x = 1$ ($T_{SG} =$

$\frac{J}{\tanh^{-1}(\frac{1}{\sqrt{\gamma}})}$). In the high temperature limit, $x < 1$ this ratio dies in the thermodynamic limit, signalling the absence of memory as discussed previously. In the low temperature regime, $x > 1$, the ratio begins to grow, diverging in the strict zero temperature limit. Therefore, this quantity $\langle \sigma_0 M_n \rangle / \Omega$ provides a good order parameter on the Cayley tree. Fig.2(b) obtains this transition temperature by studying the quantity Ω^2 scaled appropriately. Appendix C has a discussion of why this quantity is more convenient for finite-sized systems. The reason the second transition temperature is called 'spin-glass' is explained in Appendix D. Quite remarkably, the methods of Eggarter and others carry right through when one considers a $\pm J$ disordered Hamiltonian on the Cayley tree, and there occurs only one transition temperature at precisely $\frac{J}{\tanh^{-1}(\frac{1}{\sqrt{\gamma}})}$, thus justifying the name 'spin-glass'. However, look at Appendix D for a discussion on how there is not a regular type of 'spin-glass' phase below T_{SG}.

One could expect that higher moments of $\sigma_0 M_n$ would give us new transition temperatures, which would presumably be related to the infinite set of critical temperatures found for this system in [5]. With the aid of the new recursive techniques discussed in Appendix B, we were able to compute analytically the next moment, without finding any new critical temperature. We leave this interesting connection for future study.

We note that the techniques of Ref.[12] can be used in conjunction with the above insights to study the classical Heisenberg ferromagnet on a Cayley Tree. For this case, the function that we have defined as $a = \tanh \beta J$ would merely change to $a_{Heisenberg} \equiv \coth \beta J - \frac{1}{\beta J}$, and thus the two transition temperatures would be given by $\gamma a_{Heisenberg} = 1$, and $\gamma a^2_{Heisenberg} = 1$, closely analogous to the Ising case. Indeed the same logic should hold for arbitrary m-component vectors though a closed form solution is unlikely.

5. Conclusions and Outlook

We have shown that the dynamic time Markov chain approach gives exactly the same results as the static ensemble approach of statistical mechanics, thus linking two different communities. We have obtained the two transition temperatures present in this model with new recursive techniques, which may be interesting for other tree like models.

We have highlighted that $\sigma_0 M_n$ is a convenient order parameter for the system, which can at the same time be thought of as a kind of memory in the time-picture. This quantity signals the Bethe-Peirels transition temperature T_{BP}. Furthermore, we have shown that $\frac{\langle \sigma_0 M_n \rangle}{\Omega}$, where $\Omega = \sqrt{\langle (\sigma_0 M_n)^2 \rangle - \langle \sigma_0 M_n \rangle^2}$ is the RMS deviation, is the order parameter for T_{SG}. The analytical understanding allows us to tune our Monte Carlo simulations, which are usually problematic in these kinds of lattices due to the difficulties appearing when taking the thermodynamic limit, to study the right quantities showing clearly the transition temperatures.

The analogue to $\sigma_0 M_n$ in other expanding structures such Euclidean Anti de Sitter or de Sitter space-times is a very natural and known quantity, the study of which in this light we postpone for the future. Indeed, the fact that in the Cayley tree simple correlators provide the same physical results as more involved memory type questions concerning the mutual information explains the relation signalled in Ref.[9] between quantization schemes and the persistence of memory. The rate of falloff of the correlators, determined in our case by the temperature and in the EAdS and dS cases by the dimension of the operator being examined, plays the central role in this problem.

It would also be interesting to study the correlations between the central node and the sum on the $n^{th} layer$ in quantum models; the entanglement between the central node and the sum on the $n^{th} layer$ could be another intriguing direction to pursue.

Acknowledgments

A.S. thanks Arti Garg, and Peter Young for many helpful discussions and for comments on an earlier version of the paper. J.M.M wishes to thank J.L.F. Barbón for criticisms and correspondence. We are grateful to Markus Müller, Vladimir Gritsev and Dionys Baeriswyl whose observations initiated our Appendix D. Also both authors gratefully acknowledge the people of Brazil, who, through the CNPQ institution, provide financial support to our research.

Appendix A. Obtaining the order-parameters

Suppose we want to measure how correlated two random variables A, B are. The quantity that is commonly considered in the Markov process community is mutual information, which is defined as

$$I(A,B) = S(A) + S(B) - S(A,B), \tag{A.1}$$

where

$$\begin{aligned} S(A) &= -\sum_A P(A)\log(P(A)) \\ S(B) &= -\sum_B P(B)\log(P(B)) \quad \textit{and} \\ S(A,B) &= -\sum_{A,B} P(A,B)\log(P(A,B)) \end{aligned} \tag{A.2}$$

are the usual Shannon entropies. This quantity is bounded by zero from below (This happens when $P(A,B) = P(A)P(B)$, which implies that the correlator of the product factorizes $\langle AB\rangle = \langle A\rangle\langle B\rangle$). From above, it is bounded by the entropy of the random variable with lower entropy. This happens when knowing one of them implies knowing the other(for this to happen we need that the entropy stored in the first is at least as much as the entropy stored in the second). For this case, one would expect that the quantity $\langle AB\rangle - \langle A\rangle\langle B\rangle$, which is just the often-studied correlation function in statistical physics, becomes large. However, the *ratio* $\frac{\langle AB\rangle - \langle A\rangle\langle B\rangle}{\sigma_{AB}}$, where σ_{AB} is the standard deviation, must also be studied because we may have a scenario where σ_{AB} is also large, and thus the knowledge of one of the two random variables A leads to very little predictive power for the variable B even though the two have a large correlation $\langle AB \rangle$. These two different types of quantities gives us order-parameters that mark precisely the two different phase transitions discussed in the main paper.

Appendix B. Recursion Relations to compute deviation

Here we obtain the sum $\sum_{i<j} a^{d_{ij}}$, where all i, j are points on the n^{th}-layer boundary, with the help of a recursion relation that is obtained by means of a graphical iteration which constructs the tree step by step, from the boundary to the root. We begin by considering two spins at the boundary at distance 2 in tree units (the minimal distance). At this initial step the sum is, since there are $\binom{\gamma}{2}$ ways of choosing the two spins, just trivially $\binom{\gamma}{2}a^2$ and which provides an initial condition for the iteration. Now we go one step higher, by stitching together γ of these 1-layer trees to one spin above, and thus also connecting spins which are at distance 4. The sum in this case is found to be γ times the first sum, corresponding to the γ cases in which the two spins lie within the same old branch, and $\binom{\gamma}{2}\gamma^2$ terms with value equal to a^4, which give us the cases in which one spin lies in one old branch and the second spin in another. This pattern is easily seen to hold until one step before the last iteration (the last iteration should include one branch more in the tree we are studying, and has to be treated separately).

Calling this iteration S_k the recursion relation takes the following form:

$$S_{k+1} = \gamma S_k + \binom{\gamma}{2}\gamma^{2k} a^{2(k+1)}, \tag{B.1}$$

where $S_1 = \binom{\gamma}{2}a^2$. This can be seen to be a geometric sum by the following transformation $S_k \to \binom{\gamma}{2}\gamma^{k-2}C_k$. The solution is then

$$S_k = \gamma^{k-2}\frac{x^{k+1} - x}{x - 1}, \tag{B.2}$$

where $x = \gamma a^2$.

Adding $\gamma + 1$ braches to form the top root-node (because of the type of Cayley tree we have considered), we have

$$\sum_{i<j} a^{d_{ij}} = N_n\left(\frac{(\gamma - 1)}{2\gamma}\frac{x^{n+1} - x}{x - 1} + \frac{x^n}{2}\right), \tag{B.3}$$

where $N_n = (\gamma + 1)\gamma^{n-1}$ is the number of spins in the last layer.

Appendix C. Monte Carlo Simulation

Here we show how to exploit the analytical understanding to obtain the transition temperatures in the thermodynamic limit, with the help of Monte-Carlo simulation on finite-sized systems. We showed in the main paper that

$$\langle \sigma_0 M_n \rangle = (\gamma \tanh(\beta J))^n \left(1 + \frac{1}{\gamma}\right) \tag{C.1}$$

n	N	N_{run}	N_{sweep}	T_{min}	T_{max}	N_T
2	10	100	2048	0.10	5.00	50
3	22	100	2048	0.10	5.00	50
4	46	100	2048	0.10	5.00	50
5	94	100	2048	0.10	5.00	50
6	190	100	4096	0.10	5.00	50
7	382	100	4096	0.10	5.00	50
8	766	100	8192	0.30	3.00	50

Table C1. Parameters of the simulations. Here n is the number of layers in the Cayley tree, N is the number of spins (the coordination number was fixed at $z = 3$), N_{run} is the number of runs averaged over to extract error-bars, N_{sweep} is the number of Metropolis Monte Carlo sweeps, T_{min} and T_{max} are the lowest and highest temperatures simulated, and N_T is the number of temperatures used for parallel-tempering.

and argued that as $n \to \infty$, this system undergoes a sharp change in behavior at the critical temperature given by $\gamma \tanh(\beta J) = 1$. Now we notice that in fact, this quantity can show the transition even when n is finite. The key point is that at precisely the transition temperature, the quantity $(\gamma \tanh(\beta J))^n$ becomes independent of n, therefore by plotting this quantity as a function of temperature for various system-sizes on the same graph, we can instantly obtain the transition temperature as the point at which the various curves take the same value. This is done in the main paper. Next we showed in the main paper that the quantity $\frac{\langle \sigma_0 M_n \rangle}{\Omega}$ where $\Omega = \sqrt{\langle (\sigma_0 M_n)^2 \rangle - \langle \sigma_0 M_n \rangle^2}$ is the RMS deviation, has a sharp change at the spin-glass transition temperature given by $x \equiv \gamma \tanh^2(\beta J) = 1$ in the thermodynamic limit, $n \to \infty$. Now, we point out that the mean square deviation Ω^2 itself can be used in a finite-size-scaling study to obtain T_{SG}. We showed in the paper, based on the techniques described in the last section above, that,

$$\Omega^2 = \langle (\sigma_0 M_n)^2 \rangle - \langle \sigma_0 M_n \rangle^2 = N_n \left(1 + \frac{\gamma - 1}{\gamma} \frac{x^{n+1} - x}{x - 1} - \frac{1}{\gamma} x^n\right). \tag{C.2}$$

For finite n, it is better to write this as

$$\Omega^2 = N_n \left(1 + \frac{\gamma - 1}{\gamma} x (1 + x + x^2 + \cdots x^{n-1}) - \frac{1}{\gamma} x^n\right). \tag{C.3}$$

It is clear that when $x = 1$, $\Omega^2 = N_n n \frac{\gamma - 1}{\gamma}$. Therefore, the quantity $\frac{\Omega^2}{N_n n}$ becomes independent of system-size at this particular value of the temperature; thus plotting this quantity as a function of temperature for various sizes on the same graph, we are able to obtain the spin-glass transition temperature T_{SG}, as shown in the main paper.

Our simulation used the standard Metropolis algorithm in conjunction with parallel-tempering which helps speed up the equilibration. Table C1 collects the parameters of the simulations. Equilibration was verified by keeping track of the energy; those points (at very low-temperature, and large system sizes) for which the energy did not stay the same within error-bars over the last two sweeps, were discarded.

n	N	N_{samp}	N_{sweep}	T_{min}	T_{max}	N_T
2	10	100	2048	0.30	3.00	40
3	22	100	2048	0.30	3.00	40
4	46	100	2048	0.30	3.00	40
5	94	100	2048	0.30	3.00	40
6	190	100	4096	0.30	3.00	40
7	382	100	4096	0.30	3.00	40
8	766	100	8192	0.30	3.00	40

Table D1. Parameters of the simulations on $\pm J$ spin glass. Here n is the number of layers in the Cayley tree, N is the number of spins (the coordination number was fixed at $z = 3$), N_{samp} is the number of samples averaged over to extract error-bars, N_{sweep} is the number of Metropolis Monte Carlo sweeps, T_{min} and T_{max} are the lowest and highest temperatures simulated, and N_T is the number of temperatures used for parallel-tempering.

Appendix D. The Spin-Glass Transition Temperature

Here we consider the $\pm J$ spin glass on the Cayley tree given by the Hamiltonian

$$\mathcal{H} = \sum_{\langle i,j\rangle} J_{ij}\sigma_i\sigma_j, \tag{D.1}$$

where J_{ij} are independent identically distributed random variables that can take values $\pm J$ with equal probability, and where the sum involves pairs of spins which are adjacent on the tree. Instead of the spin variables, like before, we choose the spin at the root of the tree σ_0, and the bonds $\theta_{ij} = \sigma_i\sigma_j$ to define the configuration of our system. The θ variables can take values ± 1, which make them effectively spin variables too. The Hamiltonian then takes the following simple form

$$\mathcal{H} = \sum_{\langle i,j\rangle} J_{ij}\theta_{ij}. \tag{D.2}$$

The partition function for this model is identical to that for the Ising ferromagnet since it is an even function in J, but more importantly, with the help of the same trick of inserting squares of all the spins that lie between any two given spins, we can now write down the correlator:

$$\langle\sigma_i\sigma_j\rangle = \pm\tanh(\beta J)^{d_{ij}} = \pm a^{d_{ij}}. \tag{D.3}$$

Whether a plus sign appears or a minus sign appears depends on the number of bonds between i and j that are ferromagnetic. We will now compute the 'Edwards-Anderson' type order-parameter with the knowledge that memory is the order-parameter for the ferromagnet, i.e we are interested in

$$q = [\langle\sigma_0 M_n\rangle^2]_{avg}, \tag{D.4}$$

where $[\cdots]_{avg}$ means an average over disorder.



Figure D1. Monte Carlo simulation of the $\pm J$ spin glass no a Cayley trees with coordination number $z = 3$. The figure obtains the transition temperature popularly called Spin-glass T_{SG} [10]. While this temperature is of importance also in the regular ferromagnet as shown in the main paper, the reason for its name comes from studying the spin-glass as shown here.

Computing the correlation between the root spin and the sum of the spins at the n^{th} level we obtain:

$$\langle\sigma_0 M_n\rangle = \sum_{i\in n^{th} layer} \langle\sigma_0\sigma_i\rangle = \tanh(\beta J)^n \sum_{N_n terms} \pm 1, \tag{D.5}$$

where N_n is equal to the number of spins in the n^{th} layer: $N_n = z(z-1)^{n-1}$, where z is the coordination number for the type of Cayley tree shown in Fig.1. Squaring we have:

$$\langle\sigma_0 M_n\rangle^2 = \tanh(\beta J)^{2n}(\sum_{N_n terms} \pm 1)(\sum_{N_n terms} \pm 1) \tag{D.6}$$

The ± 1 terms in the two sums are completely random, so when we square and average over disorder, only the 'diagonal' terms which give $(\pm 1)^2 = 1$ stay, all other terms vanishing because $+1$ is as likely as -1. Therefore, defining $\gamma \equiv z - 1$, we have

$$[\langle\sigma_0 M_n\rangle^2]_{avg} = (\gamma\tanh(\beta J)^2)^n(1 + \frac{1}{\gamma}) \tag{D.7}$$

It is clear that the behaviour of $[\langle\sigma_0 M_n\rangle^2]$ for 'long times' or, equivalently, in the 'thermodynamic limit' $n \to \infty$, changes completely depending on whether $\gamma\tanh(\beta J)^2$ is $>$ or $<$ 1. So, $\gamma\tanh(\beta J)^2 = 1$ signals a transition which is understood to be the so-called spin-glass critical temperature $T_{SG} = \frac{J}{\tanh^{-1}(\frac{1}{\sqrt{\gamma}})}$. Fig. D1 shows that this phase transition can be obtained from Monte Carlo simulations of finite-systems. Again we exploit the convenient fact that at T_{SG}, the quantity $(\gamma\tanh(\beta J)^2)^n$ becomes independent of n, therefore by plotting q as a function of temperature for various system-sizes on the same graph, we can instantly obtain the transition temperature as the point at which the various curves take the same value. Our simulation once again used the standard Metropolis algorithm in conjunction with parallel-tempering which helps speed up the equilibration. Table D1 collects the parameters of the simulations. We make the observation here that the phase below T_{SG} is not really a 'spin-glass' in the sense that there is no complex free-energy landscape. Two essential characteristics of a spin glass are frustration and disorder; here, we have only disorder, no frustration. Thus the transition T_{SG} is more like a regular ferromagnetic phase transition.

[1] T.P.Eggarter. Cayley trees, the ising problem, and the thermodynamic limit. *Phys.Rev.B*, 9:2989–2992, Apr 1974.

[2] Hirotsugu Matsuda. Infinite susceptibility with spontaneous magnetization - exact properties of the ising model on the cayley tree. *PTP*, 51(4):1053–1063, Oct 1974.

[3] Tohru Morita and Tsuyoshi Horiguchi. Susceptibility and correlation function of the ising model on the cayley tree. *PTP*, 54(4):982–998, 1975.

[4] J von Heimburg and H Thomas. Phase transition of the cayley tree with ising interaction. *J.Phys.C*, 7(19):3433, Oct 1974.

[5] E.Müller-Hartmann and J.Zittartz. New type of phase transition. *Phys.Rev.Lett*, 33:893–897, Oct 1974.

[6] William Evans, Claire Kenyon, Yuval Peres, and Leonard J.Schulman. Broadcasting on trees and the ising model. *The Annals of Applied Probability*, 10(2):410–433, 2000.

[7] Daniel Harlow, Stephen H.Shenker, Douglas Stanford, and Leonard Susskind. Tree-like structure of eternal inflation: A solvable model. *Phys.Rev.D*, 85:063516, March 2012.

[8] Javier M. Magán José L. F. Barbón. Fast scramblers, horizons and expander graphs. *JHEP*, 16, August 2012.

[9] Daniel A.Roberts and Douglas Stanford. On memory in exponentially expanding spaces. *arXiv*, Oct 2012.

[10] Marc Mézard and Giorgio Parisi. The bethe lattice spin glass revisited. *The European Physical Journal B-Condensed Matter and Complex Systems*, 20(2):217–233, 2001.

[11] M. Ostilli. Cayley trees and bethe lattices: A concise analysis for mathematicians and physicists. *Physica A*, 391:3417–3423, June 2012.

[12] Michael E Fisher. Magnetism in one-dimensional systemsthe heisenberg model for infinite spin. *American Journal of Physics*, 32:343, 1964.